

# Facial Soft Tissue Fillers

## Facial Filler Injection Consent

This is an informed-consent document prepared to help inform you regarding the injection of filler products including, but not limited to, Juvederm Ultra, Juvederm Ultra Plus XC, Juvederm Voluma, Belotero, Restylane, Radiesse, Sculptra and other facial filler injections, including a discussion of their risks, benefits, expectations and alternative treatments. Soft tissue fillers are temporary and the duration of the change depends on the location of the injection and the type of filler used.

### PROCEDURE

Hyaluronic acid and calcium hydroxyapatite based facial filler injections are administered via a syringe, or injection, into the area(s) of the face where the patient seeks the improvement of wrinkles and folds or increased facial volume. Dermal filler is a liquid or gel injected into the tissue of the face.

A topical numbing cream can be used to reduce the discomfort of the injection(s). The treatment site is washed, and the depth of the injection(s) depends on the depth and location of the wrinkle(s) and fold(s). Additional treatments may be necessary until the area is corrected to the desired level. Photographs may be taken before, during and after the facial filler procedure as part of health information and will be used by the provider in accordance with separately signed photographic consent or denial forms.

### DISCOMFORT

The area of or around the injection site may become bruised or swollen. Pain, itching, discoloration and tenderness may also occur. Increased bleeding or bruising could also occur if you are currently taking medication or substances such as aspirin or other non-steroidal anti-inflammatory drugs such as Advil. Lumps and bumps under the skin are common. These symptoms usually resolve over a period of days or up to two weeks.

As with all injections there is a risk of infection. The syringe is sterile and standard precautions associated with injectable materials have been taken. Dermal fillers should not be used by persons who are hypersensitive to pain, have severe allergies and active inflammation or infections (i.e. cyst, pimples, rashes or hives) in the area chosen for injection. Laser treatment, chemical skin peeling or any other skin based procedures after facial filler injections carry the risk of inflammatory reaction at the treated site. Minimize sun exposure (UV/UR) and extremely cold weather until initial swelling and/or redness has gone away.

I understand that this procedure is cosmetic and that payment is my responsibility.

I have had an opportunity to fully inquire about the risks and benefits of this procedure and its alternatives. All my questions were answered to my

satisfaction and I consent to the procedure.

*I consent to the above.*

---

Witness Signature

01/12/2021

---

Patient / Agent / Guardian Signature

01/12/2021